



06-28-00

A BOX
SER

CERTIFICATION UNDER 37 CFR 1.10

I hereby certify that this New Application Transmittal and the documents referred to as enclosed therein are being deposited with the United States Postal Service on this date June 27, 2000 in an envelope as "Express Mail Post Office to Addressee" Mailing Label Number EL261868223US addressed to the: Commissioner for Patents, Washington, D.C. 20231.

Pamela Johnston
(Print Name)

Pamela Johnston
(Signature)



Commissioner for Patents
BOX PATENT APPLICATION
Washington, D.C. 20231

Hoffmann-La Roche Inc.
340 Kingsland Street
Nutley, NJ 07110
Case Docket 1097
June 27, 2000

Sir:


Transmitted herewith for filing under 37 C.F.R. § 1.53(b) is the patent application of

Inventor(s): Pascal Sebastian Bailon

For: **ERYTHROPOIETIN CONJUGATES**

Enclosed are:

1. X 3 sheet(s) of drawing. [] formal [X] informal
2. X 4 page(s) of Declaration and Power of Attorney (unsigned)
3. X 3 page(s) of Sequence Listing
4. X 1 computer disk(s) containing Sequence Listing
5. X Statement under 37 CFR § 1.821 or 37 C.F.R. § 1.825
6. X 38 pgs. of specification, 9 pgs. of claims, 1pg. of abstract



09604938-062700

7. The fee has been calculated as shown below:

CLAIMS				
FOR	NO. FILED	NO. EXTRA	RATE	FEE
TOTAL CLAIMS	60 - 20	40	x \$18	720
INDEP. CLAIMS	2 - 3	0	x \$78	--
MULTIPLE DEPENDENT CLAIMS PRESENTED				+ \$260
BASIC FEE				\$690
TOTAL				\$1410

8. X Please charge my Deposit Account No. 08-2525 in the amount of \$ 1410 . This sheet is provided in triplicate.
9. A check in the amount of \$ to cover the filing fee is enclosed.
10. X The Commissioner is hereby authorized to charge payment of the following fees or any additional fees associated with this communication or credit any overpayment to Deposit Account No. 08-2525. This sheet is provided in triplicate.
- X Any filing fees required under 37 C.F.R. § 1.16.
- X Any patent application processing fees under 37 C.F.R. § 1.17.

11. Priority - 35 U.S.C. § 119

FOREIGN PRIORITY

[] Foreign Priority of application(s) number filed on in
 is claimed under 35 U.S.C. § 119(a)-(d) or 35 U.S.C. § 365(a)-(b).

[] The certified copy(ies) has(have) been filed in prior U.S. patent application Serial No. on .

[] The certified copy(ies) will follow.

PRIORITY TO PROVISIONAL APPLICATION(S) UNDER 35 U.S.C. § 119(e)

[X] Priority of prior provisional application(s) Serial No. 60/142,254, filed on July 2, 1999, Serial No. 60/150,225, filed on August 23, 1999, Serial No. 60/151,548, filed August 31, 1999 and Serial No. 60/166,151, filed November 17, 1999 are claimed under 35 U.S.C. § 119(e).

☐ Amend the specification by inserting, before the first line, the following sentence: -- This application claims priority under 35 U.S.C. § 119(e) of provisional application(s) Serial No. _____, filed _____. --

12. RELATION BACK UNDER 35 U.S.C. § 120

(A) ☐ Amend the specification by inserting, before the first line, the following sentence: -- This is a ☐ continuation ☐ divisional of copending application(s) ☐ Serial No. _____ filed on _____. --

(B) ☐ A copy of the oath or declaration from the prior application noted above is enclosed.


13. ☐ INCORPORATION BY REFERENCE (*USEABLE IF ITEMS 12 (A) AND (B) ARE CHECKED*) AND IF YOU ARE FILING A CONTINUATION OR DIVISIONAL UNDER 37 C.F.R. § 1.53(b). **N.B. MUST INCLUDE COPY OF PRIOR SPECIFICATION**

The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under 12. (B), is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

14. ☐ The power of attorney in prior application is to:

- a. ☐ The power appears in the original papers of the prior application.
- b. ☐ Since the power does not appear in the original papers, a copy of the power in the prior application is enclosed.
- c. ☐ Recognize as associate attorney _____.
- d. Continue to address all communications to

George W. Johnston
Hoffmann-La Roche Inc.
340 Kingsland Street
Nutley, NJ 07110



Attorney for Applicant(s)
William H. Epstein
(Reg. No. 20008)
Telephone: (973) 235-3723
Telefax: (973) 235-2363

ERYTHROPOIETIN CONJUGATES

5

CROSS-REFERENCE TO RELATED APPLICATIONS

The priority of U.S. Provisional Application No. 60/142,254, filed July 2, 1999; No. 60/150,225, filed August 23, 1999; No. 60/151,548, filed August 31, 1999; and No.
10 60/166,151, filed November 17, 1999 is claimed.

BACKGROUND OF THE INVENTION

Erythropoiesis is the production of red blood cells, which occurs to offset cell
15 destruction. Erythropoiesis is a controlled physiological mechanism that enables
sufficient red blood cells to be available for proper tissue oxygenation. Naturally
occurring human erythropoietin (hEPO) is produced in the kidney and is the humoral
plasma factor which stimulates red blood cell production (Carnot, P and Deflandre, C
(1906) C.R. Acad. Sci. 143: 432; Erslev, AJ (1953 Blood 8: 349; Reissmann, KR (1950)
20 Blood 5: 372; Jacobson, LO, Goldwasser, E, Freid, W and Plzak, LF (1957) Nature 179:
6331-4). Naturally occurring EPO stimulates the division and differentiation of
committed erythroid progenitors in the bone marrow and exerts its biological activity by
binding to receptors on erythroid precursors (Krantz, BS (1991) Blood 77: 419).

25 Erythropoietin has been manufactured biosynthetically using recombinant DNA
technology (Egrie, JC, Strickland, TW, Lane, J et al. (1986) Immunobiol. 72: 213-224)
and is the product of a cloned human EPO gene inserted into and expressed in the ovarian
tissue cells of the chinese hamster (CHO cells). The primary structure of the
predominant, fully processed form of hEPO is illustrated in SEQ ID NO:1. There are two
30 disulfide bridges between Cys⁷-Cys¹⁶¹ and Cys²⁹-Cys³³. The molecular weight of the
polypeptide chain of EPO without the sugar moieties is 18,236 Da. In the intact EPO

molecule, approximately 40% of the molecular weight are accounted for by the carbohydrate groups that glycosylate the protein at glycosylation sites on the protein (Sasaki, H, Bothner, B, Dell, A and Fukuda, M (1987) J. Biol. Chem. 262: 12059).

5 Because human erythropoietin is essential in red blood cell formation, the hormone is useful in the treatment of blood disorders characterized by low or defective red blood cell production. Clinically, EPO is used in the treatment of anemia in chronic renal failure patients (CRF) (Eschbach, JW, Egri, JC, Downing, MR et al. (1987) NEJM 316: 73-78; Eschbach, JW, Abdulhadi, MH, Browne, JK et al. (1989) Ann. Intern. Med. 111: 992; Egrie, JC, Eschbach, JW, McGuire, T, Adamson, JW (1988) Kidney Intl. 33: 262; Lim, VS, Degowin, RL, Zavala, D et al. (1989) Ann. Intern. Med. 110: 108-114) and in AIDS and cancer patients undergoing chemotherapy (Danna, RP, Rudnick, SA, Abels, RI In: MB, Garnick, ed. Erythropoietin in Clinical Applications-An International Perspective. New York, NY: Marcel Dekker; 1990: p. 301-324). However, the
10 bioavailability of commercially available protein therapeutics such as EPO is limited by their short plasma half-life and susceptibility to protease degradation. These shortcomings prevent them from attaining maximum clinical potency.

20 SUMMARY OF THE INVENTION

20 This invention provides an erythropoietin conjugate, said conjugate comprising an erythropoietin glycoprotein having at least one free amino group and having the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells and selected from the group consisting of human
25 erythropoietin and analogs thereof which have sequence of human erythropoietin modified by the addition of from 1 to 6 glycosylation sites or a rearrangement of at least one glycosylation site; said glycoprotein being covalently linked to "n" poly(ethylene glycol) groups of the formula $-\text{CO}-(\text{CH}_2)_x-(\text{OCH}_2\text{CH}_2)_m-\text{OR}$ with the $-\text{CO}$ (i.e. carbonyl) of each poly(ethylene glycol) group forming an amide bond with one of said
30 amino groups; wherein R is lower alkyl; x is 2 or 3; m is from about 450 to about 900; n is from 1 to 3; and n and m are chosen so that the molecular weight of the conjugate

minus the erythropoietin glycoprotein is from 20 kilodaltons to 100 kilodaltons. This invention further provides compositions containing conjugates described herein in which the percentage of conjugates in the composition in which n is 1 is at least ninety percent.

Compared to unmodified EPO (i.e., EPO without a PEG attached) and conventional PEG-EPO conjugates, the present conjugates have an increased circulating half-life and plasma residence time, decreased clearance, and increased clinical activity *in vivo*. The conjugates of this invention have the same uses as EPO. In particular, the conjugates of this invention are useful to treat patients by stimulating the division and differentiation of committed erythroid progenitors in the bone marrow in the same way EPO is used to treat patients.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1: Influence of pH on thermal stability. The transition temperature is plotted against the pH.

Figure 2: Influence of ionic strength on thermal stability. The transition temperature is plotted against the phosphate concentration.

Figure 3: Dependence of thermal stability on buffer substance.

Figure 4 shows that sulfate is also a suitable buffer/additive at low pH (e.g. pH 6.2), whereas phosphate is less suitable at pH 6.2 compared to pH 7.5. This shows that sulfate keeps the thermal stability high, even at low pH.

Figure 5: Dependency of peg-EPO aggregation on pH. Peg-EPO samples after heat stress (as described above) were analyzed by SDS-PAGE. Proteins were stained with silver. Lane 1: molecular weight standard. Lane 2: pH 5. Lane 3: pH 5, reduced. Lane 4: pH 6. Lane 5: pH 6, reduced. Lane 6: pH 6.5. Lane 7: pH 6.5, reduced. Lane 8: pH 7. Lane 9: pH 7, reduced. Lane 10: peg-EPO, unstressed.

Figure 6 shows that the use of 1 mg/ml acetylcysteine as an antioxidant prevents the formation of aggregates under heat stress.

DETAILED DESCRIPTION OF THE INVENTION

5

This invention provides conjugates, said conjugates comprising an erythropoietin glycoprotein having at least one free amino group and having the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells and selected from the group consisting of human erythropoietin and
10 analogs thereof which have sequence of human erythropoietin modified by the addition of from 1 to 6 glycosylation sites or a rearrangement of at least one glycosylation site; said glycoprotein being covalently linked to "n" poly(ethylene glycol) groups of the formula $-\text{CO}-(\text{CH}_2)_x-(\text{OCH}_2\text{CH}_2)_m-\text{OR}$ with the $-\text{CO}$ (i.e. carbonyl) of each poly(ethylene glycol) group forming an amide bond with one of said amino groups;
15 wherein R is lower alkyl; x is 2 or 3; m is from about 450 to about 900; n is from 1 to 3; and n and m are chosen so that the molecular weight of the conjugate minus the erythropoietin glycoprotein is from 20 kilodaltons to 100 kilodaltons.

It has been found that the conjugates of this invention can be used in the same
20 manner as unmodified EPO. However, the conjugates of this invention have an increased circulating half-life and plasma residence time, decreased clearance, and increased clinical activity *in vivo*. Because of these improved properties, the conjugates of this invention can be administered once weekly instead of the three times weekly for unmodified EPO. Decreased frequency of administration is expected to result in
25 improved patient compliance leading to improved treatment outcomes, as well as improved patient quality of life. Compared to conventional conjugates of EPO linked to poly(ethylene glycol) it has been found that conjugates having the molecular weight and linker structure of the conjugates of this invention have an improved potency, stability, AUC, circulating half-life, and cost of goods profile.


30

The conjugates in accordance of this invention can be administered in a therapeutically effective amount to patients in the same way EPO is administered. The therapeutically effective amount is that amount of conjugate necessary for the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells. The exact amount of conjugate is a matter of preference subject to such factors as the exact type of condition being treated, the condition of the patient being treated, as well as the other ingredients in the composition. The pharmaceutical compositions containing the conjugate may be formulated at a strength effective for administration by various means to a human patient experiencing blood disorders characterized by low or defective red blood cell production. Average therapeutically effective amounts of the conjugate may vary and in particular should be based upon the recommendations and prescription of a qualified physician.

The erythropoietin glycoprotein products prepared in accordance with this invention may be prepared in pharmaceutical compositions suitable for injection with a pharmaceutically acceptable carrier or vehicle by methods known in the art. Among the preferred pharmaceutically acceptable carriers for formulating the products of the invention are human serum albumin, human plasma proteins, etc.

The term "erythropoietin" or "EPO" refers to a glycoprotein, having the amino acid sequence set out in (SEQ ID NO: 1) or (SEQ ID NO: 2) or an amino acid sequence substantially homologous thereto, whose biological properties relate to the stimulation of red blood cell production and the stimulation of the division and differentiation of committed erythroid progenitors in the bone marrow. As used herein, these terms include such proteins modified deliberately, as for example, by site directed mutagenesis or accidentally through mutations. These terms also include analogs having from 1 to 6 additional sites for glycosylation, analogs having at least one additional amino acid at the carboxy terminal end of the glycoprotein, wherein the additional amino acid includes at least one glycosylation site, and analogs having an amino acid sequence which includes a rearrangement of at least one site for glycosylation. These terms include both natural and recombinantly produced human erythropoietin.

The erythropoietin conjugates of this invention can be represented by Formula 1:



5 wherein x, m, n and R are as above. In Formula I, P is the residue of an erythropoietin glycoprotein described herein, (i.e. without the amino group or amino groups which form an amide linkage with the carbonyl shown in Formula I), having the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells. P may be selected from the group consisting of residues of human
10 erythropoietin and analogs thereof having from 1 to 6 additional sites for glycosylation. As set out in detail below, the preparation and purification of EPO are well known in the art. By EPO is meant the natural or recombinant protein, preferably human, as obtained from any conventional source such as tissues, protein synthesis, cell culture with natural or recombinant cells. Any protein having the activity of EPO, such as muteins or
15 otherwise modified proteins, is encompassed. Recombinant EPO may be prepared via expression in CHO-, BHK- or HeLa cell lines, by recombinant DNA technology or by endogenous gene activation. Expression of proteins, including EPO, by endogenous gene activation is well known in the art and is disclosed, for example in U.S. Patents Nos. 5,733,761, 5,641,670, and 5,733,746, and international patent publication Nos. WO
20 93/09222, WO 94/12650, WO 95/31560, WO 90/11354, WO 91/06667 and WO 91/09955, the contents of each of which are incorporated herein by reference. The preferred EPO species for the preparation of erythropoietin glycoprotein products are human EPO species. More preferably, the EPO species is the human EPO having the amino acid sequence set out in SEQ ID NO:1 or SEQ ID NO:2, more preferably the
25 amino acid sequence SEQ ID NO:1.

In an embodiment, P may be the residue of a glycoprotein analog having from 1 to 6 additional sites for glycosylation. Glycosylation of a protein, with one or more oligosaccharide groups, occurs at specific locations along a polypeptide backbone and
30 greatly affects the physical properties of the protein such as protein stability, secretion, subcellular localization, and biological activity. Glycosylation is usually of two types.

O-linked oligosaccharides are attached to serine or threonine residues and N-linked oligosaccharides are attached to asparagine residues. One type of oligosaccharide found on both N-linked and O-linked oligosaccharides is N-acetylneuraminic acid (sialic acid), which is a family of amino sugars containing 9 or more carbon atoms. Sialic acid is usually the terminal residue on both N-linked and O-linked oligosaccharides and, because it bears a negative charge, confers acidic properties to the glycoprotein. Human erythropoietin, having 165 amino acids, contains three N-linked and one O-linked oligosaccharide chains which comprise about 40% of the total molecular weight of the glycoprotein. N-linked glycosylation occurs at asparagine residues located at positions 24, 38, and 83 and O-linked glycosylation occurs at a serine residue located at position 126. The oligosaccharide chains are modified with terminal sialic acid residues. Enzymatic removal of all sialic acid residues from the glycosylated erythropoietin results in loss of *in vivo* activity but not *in vitro* activity because sialylation of erythropoietin prevents its binding, and subsequent clearance, by hepatic binding protein.

The glycoproteins of the present invention include analogs of human erythropoietin with one or more changes in the amino acid sequence of human erythropoietin which result in an increase in the number of sites for sialic acid attachment. These glycoprotein analogs may be generated by site-directed mutagenesis having additions, deletions, or substitutions of amino acid residues that increase or alter sites that are available for glycosylation. Glycoprotein analogs having levels of sialic acid greater than those found in human erythropoietin are generated by adding glycosylation sites which do not perturb the secondary or tertiary conformation required for biological activity. The glycoproteins of the present invention also include analogs having increased levels of carbohydrate attachment at a glycosylation site which usually involve the substitution of one or more amino acids in close proximity to an N-linked or O-linked site. The glycoproteins of the present invention also include analogs having one or more amino acids extending from the carboxy terminal end of erythropoietin and providing at least one additional carbohydrate site. The glycoproteins of the present invention also include analogs having an amino acid sequence which includes a rearrangement of at least one site for glycosylation. Such a rearrangement of glycosylation site involves the deletion of one or

more glycosylation sites in human erythropoietin and the addition of one or more non-naturally occurring glycosylation sites. Increasing the number of carbohydrate chains on erythropoietin, and therefore the number of sialic acids per erythropoietin molecules may confer advantageous properties such as increased solubility, greater resistance to proteolysis, reduced immunogenicity, increased serum half-life, and increased biological activity. Erythropoietin analogs with additional glycosylation sites are disclosed in more detail in European Patent Application 640 619, to Elliot published March 1, 1995.

In a preferred embodiment, the glycoproteins of the present invention comprise an amino acid sequence which includes at least one additional site for glycosylation such as, but not limited to, erythropoietins comprising the sequence of human erythropoietin modified by a modification selected from the following:

Asn³⁰Thr³²;

Asn⁵¹Thr⁵³;

Asn⁵⁷Thr⁵⁹;

Asn⁶⁹;

Asn⁶⁹Thr⁷¹;

Ser⁶⁸Asn⁶⁹Thr⁷¹;

Val⁸⁷Asn⁸⁸Thr⁹⁰;

Ser⁸⁷Asn⁸⁸Thr⁹⁰;

Ser⁸⁷Asn⁸⁸Gly⁸⁹Thr⁹⁰;

Ser⁸⁷Asn⁸⁸Thr⁹⁰Thr⁹²;

Ser⁸⁷Asn⁸⁸Thr⁹⁰Ala¹⁶²;

Asn⁶⁹Thr⁷¹Ser⁸⁷Asn⁸⁸Thr⁹⁰;

Asn³⁰Thr³²Val⁸⁷Asn⁸⁸Thr⁹⁰;

Asn⁸⁹Ile⁹⁰Thr⁹¹;

Ser⁸⁷Asn⁸⁹Ile⁹⁰Thr⁹¹;

Asn¹³⁶Thr¹³⁸;

Asn¹³⁸Thr¹⁴⁰;

Thr¹²⁵; and

Pro¹²⁴Thr¹²⁵.

The notation used herein for modification of amino acid sequence means that the position(s) of the corresponding unmodified protein (e.g. hEPO of SEQ ID NO:1 or SEQ ID NO:2) indicated by the superscripted number(s) is changed to the amino acid(s) that immediately precede the respective superscripted number(s).

5

The glycoprotein may also be an analog having at least one additional amino acid at the carboxy terminal end of the glycoprotein, wherein the additional amino acid includes at least one glycosylation site. The additional amino acid may comprise a peptide fragment derived from the carboxy terminal end of human chorionic gonadotropin. Preferably, the glycoprotein is an analog selected from the group consisting of (a) human erythropoietin having the amino acid sequence, Ser Ser Ser Ser Lys Ala Pro Pro Pro Ser Leu Pro Ser Pro Ser Arg Leu Pro Gly Pro Ser Asp Thr Pro Ile Leu Pro Gln (SEQ ID NO:3), extending from the carboxy terminus; (b) the analog in (a) further comprising Ser⁸⁷ Asn⁸⁸ Thr⁹⁰ EPO; and (c) the analog in (a) further comprising Asn³⁰ Thr³² Val⁸⁷ Asn⁸⁸ Thr⁹⁰ EPO.

The glycoprotein may also be an analog having an amino acid sequence which includes a rearrangement of at least one site for glycosylation. The rearrangement may comprise a deletion of any of the N-linked carbohydrate sites in human erythropoietin and an addition of an N-linked carbohydrate site at position 88 of the amino acid sequence of human erythropoietin. Preferably, the glycoprotein is an analog selected from the group consisting of Gln²⁴ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰ EPO; Gln³⁸ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰ EPO; and Gln⁸³ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰ EPO.

As used herein, "lower alkyl" means a linear or branched alkyl group having from one to six carbon atoms. Examples of lower alkyl groups include methyl, ethyl and isopropyl. In accordance with this invention, R is any lower alkyl. Conjugates in which R is methyl are preferred.

The symbol "m" represents the number of ethylene oxide residues (OCH₂CH₂) in the poly(ethylene oxide) group. A single PEG subunit of ethylene oxide has a molecular

alkoxy-PEG-SBA). Compounds of Formula II in which x is 2 are alpha-lower alkoxy, propionic acid succinimidyl esters of poly(ethylene glycol) (lower alkoxy-PEG-SPA). Any conventional method of reacting an activated ester with an amine to form an amide can be utilized. In the reaction described above, the exemplified succinimidyl ester is a leaving group causing the amide formation. The use of succinimidyl esters such as the compounds of formula II to produce conjugates with proteins are disclosed in U.S. Patent No. 5,672,662, issued September 30, 1997 (Harris, et al.).

Human EPO contains nine free amino groups, the amino-terminal amino group plus the ϵ -amino groups of 8 lysine residues. When the pegylation reagent was combined with a SBA compound of Formula II, it has been found that at pH 7.5, a protein:PEG ratio of 1:3, and a reaction temperature of from 20-25°C, a mixture of mono-, di-, and trace amounts of the tri-pegylated species were produced. When the pegylation reagent was a SPA compound of Formula II, at similar conditions except that the protein:PEG ratio was 1:2, primarily the mono-pegylated species is produced. The pegylated EPO can be administered as a mixture, or as the cation exchange chromatography separated different pegylated species. By manipulating the reaction conditions (e.g., ratio of reagents, pH, temperature, protein concentration, time of reaction etc.), the relative amounts of the different pegylated species can be varied.

Human erythropoietin (EPO) is a human glycoprotein which stimulates the formation of erythrocytes. Its preparation and therapeutic application are described in detail for example in U.S. Patent Nos. 5,547,933 and 5,621,080, EP-B 0 148 605, Huang, S.L., Proc. Natl. Acad. Sci. USA (1984) 2708-2712, EP-B 0 205 564, EP-B 0 209 539 and EP-B 0 411 678 as well as Lai, P.H. et al., J. Biol. Chem. 261 (1986) 3116-3121, an Sasaki, H. et al., J. Biol. Chem. 262 (1987) 12059-12076. Erythropoietin for therapeutic uses may be produced by recombinant means (EP-B 0 148 605, EP-B 0 209 539 and Egrie, J.C., Strickland, T.W., Lane, J. et al. (1986) Immunobiol. 72: 213-224).

Methods for the expression and preparation of erythropoietin in serum free medium are described for example in WO 96/35718, to Burg published 14 November 1996, and in

European Patent Publication No. 513 738, to Koch published 12 June 1992. In addition to the publications mentioned above, it is known that a serum-free fermentation of recombinant CHO cells which contain an EPO gene can be carried out. Such methods are described for example in EP-A 0 513 738, EP-A 0 267 678 and in a general form by
5 Kawamoto, T. et al., Analytical Biochem. 130 (1983) 445-453, EP-A 0 248 656, Kowar, J. and Franek, F., Methods in Enzymology 421 (1986) 277-292, Bavister, B., Expcology 271 (1981) 45-51, EP-A 0 481 791, EP-A 0 307 247, EP-A 0 343 635, WO 88/00967.

In EP-A 0 267 678 an ion exchange chromatography on S-Sepharose, a preparative reverse phase HPLC on a C₈ column and a gel filtration chromatography are described
10 for the purification of EPO produced in serum-free culture after dialysis. In this connection the gel filtration chromatography step can be replaced by ion exchange chromatography on S-Sepharose fast flow. It is also proposed that a dye chromatography on a Blue Trisacryl column be carried out before the ion exchange chromatography.

A process for the purification of recombinant EPO is described by Nobuo, I. et al., J.
15 Biochem. 107 (1990) 352-359. In this process EPO is treated however with a solution of Tween[®] 20, phenylmethylsulfonyl fluoride, ethylmaleimide, pepstatin A, copper sulfate and oxamic acid prior to the purification steps. Publications, including WO 96/35718, to Burg published 14 November 1996, discloses a process for preparing erythropoietin in a
20 serum free fermentation process (EPOsf).

The specific activity of EPO or EPO conjugates in accordance with this invention can be determined by various assays known in the art. The biological activity of the purified EPO proteins of this invention are such that administration of the EPO protein by
25 injection to human patients results in bone marrow cells increasing production of reticulocytes and red blood cells compared to non-injected or control groups of subjects. The biological activity of the EPO proteins, or fragments thereof, obtained and purified in accordance with this invention can be tested by methods according to Annable, et al., Bull. Wld. Hlth. Org. (1972) 47: 99-112 and Pharm. Europa Spec. Issue Erythropoietin
30 BRP Bio 1997(2). Another biological assay for determining the activity of EPO protein, the normocythaemic mouse assay, is described in Example 4.

002290-3764960

This invention provides a composition comprised of conjugates as described above. A composition containing at least ninety percent mono-PEG conjugates, i.e. in which n is 1, can be prepared as shown in Example 5. Usually mono-PEG conjugates of erythropoietin glycoproteins are desirable because they tend to have higher activity than di-PEG conjugates.. The percentage of mono-PEG conjugates as well as the ratio of mono- and di-PEG species can be controlled by pooling broader fractions around the elution peak to decrease the percentage of mono-PEG or narrower fractions to increase the percentage of mono-PEG in the composition. About ninety percent mono-PEG conjugates is a good balance of yield and activity. Sometimes compositions in which, for example, at least ninety-two percent or at least ninety-six percent of the conjugates are mono-PEG species (n equals 1) may be desired. In an embodiment of this invention the percentage of conjugates where n is 1 is from ninety percent to ninety-six percent.

The invention will be better understood by reference to the following examples which illustrate but do not limit the invention described herein.

EXAMPLE 1: Fermentation And Purification Of Human EPO

a) Inoculum Preparation and Fermentation

One vial of the Working Cell Bank, originating from an EPO-producing CHO cell line (ATCC CRL8695, disclosed in EP 411 678 (Genetics Institute) can be used) is taken from the gas phase of the liquid nitrogen storage tank. The cells are transferred into glass spinner flasks and cultivated in a hydrogen carbonate-buffered medium in a humidified CO₂ incubator. Typical serum free media used for the inoculum preparation and fermentation are disclosed in European Patent Application 513 738, to Koch published 12 June 1992, or WO 96/35718, to Burg published 14 November 1996, for example contain as medium DMEM/F12 (e.g. JRH Biosciences/Hazleton Biologics, Denver, US, order No. 57-736) and additionally sodium hydrogencarbonate, L+glutamine, D+glucose, recombinant insulin, sodium selenite, diaminobutane, hydrocortisone, iron(II) sulfate, asparagine, aspartic acid, serine and a stabilizer for mammalian cells such as e.g. polyvinyl alcohol, methyl cellulose, polydextran, polyethylene glycol, Pluronic F68, plasma expander polygelin (HEMACCEL[®]) or polyvinyl pyrrolidone (WO 96/35718).

The cultures are microscopically checked for the absence of contaminating microorganisms, and the cell densities are determined. These tests are performed at each splitting step.

5 After the initial growth period, the cell culture is diluted with fresh medium to the starting cell density and undergoes another growth cycle. This procedure is repeated until a culture volume of approximately 2 l per glass spinner flask has been obtained. After approx. 12 doublings 1 to 5 liter of this culture is available which then is used as inoculum for the 10 l inoculum fermenter.

10 After 3 - 5 days, the culture in the 10 l fermenter can be used as inoculum for the 100 l inoculum fermenter.

15 After additional 3 - 5 days of cultivation, the culture in the 100 l fermenter can be used as inoculum for the 1000 l production fermenter.

b) Harvesting and Cell Separation

20 A batch refeed process is used, i.e. when the desired cell density is reached, approx. 80 % of the culture is harvested. The remaining culture is replenished with fresh culture medium and cultivated until the next harvest. One production run consists of a maximum of 10 subsequent harvests: 9 partial harvests and 1 overall harvest at the end of fermentation. Harvesting takes place every 3 - 4 days.

25 The determined harvest volume is transferred into a cooled vessel. The cells are removed by centrifugation or filtration and discarded. The EPO containing supernatant of the centrifugation step is in-line filtered and collected in a second cooled vessel. Each harvest is processed separately during purification.

A typical process for the purification of EPO-protein is disclosed in WO 96/35718, to Burg published 14 November 1996. The purification process is explained in the following.

5 a) Blue Sepharose Chromatography

Blue Sepharose (Pharmacia) consists of Sepharose beads to the surface of which the Cibacron blue dye is covalently bound. Since EPO binds more strongly to Blue Sepharose than most non-proteinaceous contaminants, some proteinaceous impurities and
10 PVA, EPO can be enriched in this step. The elution of the Blue Sepharose column is performed by increasing the salt concentration as well as the pH.

The column is filled with 80 - 100 l of Blue Sepharose, regenerated with NaOH and equilibrated with equilibration buffer (sodium/ calcium chloride and sodium acetate). The
15 acidified and filtered fermenter supernatant is loaded. After completion of the loading, the column is washed first with a buffer similar to the equilibration buffer containing a higher sodium chloride concentration and consecutively with a Tris-base buffer. The product is eluted with a Tris-base buffer and collected in a single fraction in accordance with the master elution profile.

20 b) Butyl Toyopearl Chromatography

The Butyl Toyopearl 650 C (Toso Haas) is a polystyrene based matrix to which aliphatic butyl-residues are covalently coupled. Since EPO binds more strongly to this gel
25 than most of the impurities and PVA, it has to be eluted with a buffer containing isopropanol.

The column is packed with 30 - 40 l of Butyl Toyopearl 650 C, regenerated with NaOH, washed with a Tris-base buffer and equilibrated with a Tris-base buffer
30 containing isopropanol.

The Blue Sepharose eluate is adjusted to the concentration of isopropanol in the column equilibration buffer and loaded onto the column. Then the column is washed with equilibration buffer with increased isopropanol concentration. The product is eluted with elution buffer (Tris-base buffer with high isopropanol content) and collected in a single fraction in accordance with the master elution profile.

c) Hydroxyapatite Ultrogel Chromatography

The Hydroxyapatite Ultrogel (Biosepra) consists of hydroxyapatite which is incorporated in an agarose matrix to improve the mechanical properties. EPO has a low affinity to hydroxyapatite and can therefore be eluted at lower phosphate concentrations than protein impurities.

The column is filled with 30 - 40 l of Hydroxyapatite Ultrogel and regenerated with a potassium phosphate/ calcium chloride buffer and NaOH followed by a Tris-base buffer. Then it is equilibrated with a Tris-base buffer containing a low amount of isopropanol and sodium chloride.

The EPO containing eluate of the Butyl Toyopearl chromatography is loaded onto the column. Subsequently the column is washed with equilibration buffer and a Tris-base buffer without isopropanol and sodium chloride. The product is eluted with a Tris-base buffer containing a low concentration of potassium phosphate and collected in a single fraction in accordance with the master elution profile.

d) Reversed Phase HPLC on Vydac C4

The RP-HPLC material Vydac C4 (Vydac) consists of silica gel particles, the surfaces of which carry C4-alkyl chains. The separation of EPO from the proteinaceous impurities is based on differences in the strength of hydrophobic interactions. Elution is performed with an acetonitrile gradient in diluted trifluoroacetic acid.

Preparative HPLC is performed using a stainless steel column (filled with 2.8 to 3.2 liter of Vydac C4 silicagel). The Hydroxyapatite Ultrogel eluate is acidified by adding trifluoro-acetic acid and loaded onto the Vydac C4 column. For washing and elution an acetonitrile gradient in diluted trifluoroacetic acid is used. Fractions are collected and immediately neutralized with phosphate buffer. The EPO fractions which are within the IPC limits are pooled.

e) DEAE Sepharose Chromatography

The DEAE Sepharose (Pharmacia) material consists of diethylaminoethyl (DEAE) - groups which are covalently bound to the surface of Sepharose beads. The binding of EPO to the DEAE groups is mediated by ionic interactions. Acetonitrile and trifluoroacetic acid pass through the column without being retained. After these substances have been washed off, trace impurities are removed by washing the column with acetate buffer at a low pH. Then the column is washed with neutral phosphate buffer and EPO is eluted with a buffer with increased ionic strength.

The column is packed with DEAE Sepharose fast flow. The column volume is adjusted to assure an EPO load in the range of 3 - 10 mg EPO/ml gel. The column is washed with water and equilibration buffer (sodium/ potassium phosphate). The pooled fractions of the HPLC eluate are loaded and the column is washed with equilibration buffer. Then the column is washed with washing buffer (sodium acetate buffer) followed by washing with equilibration buffer. Subsequently, EPO is eluted from the column with elution buffer (sodium chloride, sodium/ potassium phosphate) and collected in a single fraction in accordance with the master elution profile.

The eluate of the DEAE Sepharose column is adjusted to the specified conductivity. The resulting drug substance is sterile filtered into Teflon bottles and stored at -70°C .

EXAMPLE 2: Pegylation of EPO with mPEG-SBA

EPO purified in accordance with the serum free procedure of Example 1 (EPOsf) was homogeneous as determined by analytical methods and showed the typical isoform pattern consisting of 8 isoforms. It had a specific biological activity of 190,000 IU/mg as determined by the normocythaemic mouse assay. The pegylation reagent used was a methoxy-PEG-SBA, which is a compound of Formula II in which R is methyl; x is 3; and m is from 650 to 750 (avg. about 680, corresponding to an average molecular weight of about 30 kDa).

Pegylation Reaction

To one hundred milligrams of EPOsf (9.71 ml of a 10.3 mg/ml EPOsf stock, 5.48 μ mol) 10 ml of 0.1 M potassium phosphate buffer, pH, 7.5 containing 506 mg of 30kDa methoxy-PEG-SBA (16.5 μ mol) (obtained from Shearwater Polymers, Inc., Huntsville, Alabama) was added and mixed for 2h at room temperature (20-23 °C). The final protein concentration was 5 mg/ml and the protein:PEG reagent ratio was 1:3. After two hours, the reaction was stopped by adjusting the pH to 4.5 with glacial acetic acid and stored at -20°C, until ready for purification.

Purification

1. **Conjugate Mixture:** Approximately 28 ml of SP-SEPHAROSE FF (sulfo-propyl cation exchange resin) was packed into an AMICON glass column (2.2 x 7.5 cm) and equilibrated with 20 mM acetate buffer pH, 4.5 at a flowrate of 150 ml/h. Six milliliters of the reaction mixture containing 30 mg protein was diluted 5-fold with the equilibration buffer and applied onto the column. Unadsorbed materials were washed away with the buffer and the adsorbed PEG conjugate mixture was eluted from the column with 0.175 M NaCl in the equilibration buffer. Unmodified EPOsf still remaining on the column was eluted with 750 mM NaCl. Column was reequilibrated in the starting buffer. Samples were analyzed by SDS-PAGE and their

degree of pegylation were determined. It was found that the 0.175M NaCl eluate contained, mono- as well as di- and trace amounts of the tri-pegylated species, whereas the 750 mM NaCl eluate contained unmodified EPOsf.

- 5 2. **Di-PEG and Mono-PEG-EPOsf:** The purified conjugate mixture eluted from the column in the previous step was diluted 4-fold with the buffer and reapplied onto the column and washed as described. Di-PEG-EPOsf and mono-PEG-EPOsf were separately eluted from the column with 0.1M NaCl and 0.175 M NaCl, respectively. Elution was also performed with 750mM NaCl to elute any remaining unmodified EPOsf.
- 10

Alternatively, the reaction mixture was diluted 5-fold with the acetate buffer and applied onto the SP-Sepharose column (~0.5 mg protein/ml gel). Column was washed and adsorbed mono-PEG-EPOsf, di-PEG-EPOsf and unmodified EPOsf were eluted as described in the previous section.

15

Results

PEG-EPOsf was synthesized by chemically conjugating a linear PEG molecule with a number average molecular weight of 30 kDa. PEG-EPOsf was derived from the reaction between the primary amino groups of EPOsf and the succinimidyl ester derivative of a 30 kDa PEG-butyric acid, resulting in an amide bond.

20

Results are summarized in Table1. Purified conjugate mixture comprised of mono- and di-PEG-EPOsf and was free of unmodified EPOsf as determined by SDS-PAGE analysis. Conjugate mixture accounted for 23.4 mg or 78% of the starting material. Cation exchange chromatographic separation of mono- and di-PEG-EPOsf indicated that mono- to di-PEG ratio in the conjugate mixture was almost 1:1. After completion of the reaction, ratio of the individual components of Mono: Di: Unmodified were 40: 38: 20 (%).

25

30 Overall yield was almost quantitative.

Table 1. Summary of results of EPOsf pegylation

	<u>Sample</u>	<u>Protein (mg)</u>	<u>Yield (%)</u>
5	Rxn. Mix.	30	100
	Mono-	12.0	40
	Di-	11.4	38
10	Unmod.	6.0	20
	Conju. Mix.	23.4	78

15

EXAMPLE 3: Pegylation of EPO with mPEG-SPA

A different aliquot of the EPOsf used in Example 2 was reacted with 30 kDa methoxy-PEG-SPA (Shearwater Polymers, Inc., Huntsville, Alabama). Reaction was performed at a protein:reagent ratio of 1:2 and purification techniques were in accordance with Example 2. Primarily the mono-pegylated species was produced.

EXAMPLE 4: In-vivo activity of pegylated EPO determined by the normocythaemic mouse assay

The normocythaemic mouse bioassay is known in the art (Pharm. Europa Spec. Issue Erythropoietin BRP Bio 1997(2)) and a method in the monography of erythropoietin of Ph. Eur. BRP. The samples were diluted with BSA-PBS. Normal healthy mice, 7-15 weeks old, were administered s.c. 0.2 ml of the EPO-fraction containing un-pegylated EPO or tri-, di- or mono-pegylated EPO from Example 2 or 3. Over a period of 6 days, blood was drawn by puncture of the tail vein and diluted such that 1 µl of blood was present in 1 ml of an 0.15 µmol acridine orange staining solution. The staining time was 3 to 10 minutes. The reticulocyte counts were carried out microfluorometrically in a flow cytometer by analysis of the red fluorescence histogram. The reticulocyte counts were

given in terms of absolute figures (per 30,000 blood cells analyzed). For the data presented, each group consisted of 5 mice per day, and the mice were bled only once.

In separate experiments, a single dose of unmodified EPO (25 ng of EPO), the PEG(SBA)-EPO mixture from Example 2 (10 ng of conjugate), mono- and di- pegylated EPOs from Example 2 (10 ng of conjugate), the PEG(SPA)-EPO from Example 3 (10 ng of conjugate), and buffer solution were administered to mice. The results are shown in Table 2. The results show the superior activity and the prolonged half life of the pegylated EPO species indicated by the significantly increased amounts of reticulocytes and the shift of the reticulocytes count maximum using the same dose per mouse (10 ng), compared to a dose of 25 ng for unmodified EPO.

TABLE 2

	EPO (Unmodified)	30 kDa SPA PEG	Mono 30K SBA	Di 30K SBA	PEG-EPO SBA Conjugate Mixture	Control Buffer
72h	1000	1393	1411	994	1328	857
96h	500	1406	1501	926	1338	697
120h	~200	1100	1182	791	944	701
144h	~0	535	607	665	660	708

5

EXAMPLE 5: Preparation of Predominantly mono-PEG-EPO**Pegylation Reaction**

10

Starting with 100 mg (5.48 μ mol) of EPOsf in 100 mM potassium phosphate buffer pH 7.5 prepared in accordance with Example 1, there was added 329 mg (10.96 μ mol) of 30 kDa PEG-SBA reagent dissolved in 3ml 1 mM HCL. Enough 100 mM potassium phosphate buffer pH 7.5 was added to make the reaction mixture volume to 20 ml. The final protein concentration was 5 mg/ml and the protein : PEG reagent ratio was 1:2. The reaction mixture was mixed for 2h at ambient temperature (20 - 22°C). After 2h, the reaction was stopped by adjusting the pH to 4.5 with glacial acetic acid and stored frozen at -20°C until ready for purification.

Purification

20

The reaction mixture from the previous step was diluted 1:5 with 10 mM sodium acetate, pH 4.5 and applied to 300 ml SP-Sepharose FF (sulfopropyl cation exchange resin) packed into a 4.2 x 19 cm column. The column was previously equilibrated with the same buffer. Column effluents were monitored at 280 nm with a Gilson UV monitor and recorded with a Kipp and Zonen recorder. The column was washed with 300 ml or 1 bed volume of equilibration buffer to remove excess reagents, reaction byproducts and

25

oligomeric PEG-EPO. It was followed by washing with 2 bed volumes of 100 mM NaCl to remove di-PEG-EPO. Mono-PEG-EPO was then eluted with 200 mM NaCl. During elution of the mono-PEG-EPO, the first 50 ml of the protein peak was discarded and the mono-PEG-EPO was collected as a 150 ml fraction. Unmodified EPOsf remaining on the column was eluted with 750 mM NaCl. All elution buffers were made in the equilibration buffer. All eluted samples were analyzed by SDS-PAGE and by high performance Size Exclusion Chromatography (SEC). The mono-PEG-EPO pool obtained from the 150 ml fraction, which had no detectable unmodified EPOsf, was then concentrated to ~ 4.5 – 7.5 mg/ml and diafiltered into the storage buffer, 10 mM potassium phosphate, 100 mM NaCl , pH 7.5. Concentration/Diafiltration was performed with Millipore Labscale™ TFF System fitted with 50 kDa cut off Millipore Pellicon XL Biomax 50 membrane at ambient temperature. Concentrated mono-PEG-EPO was sterile filtered and stored frozen at -20°C.

Approximately 75% of EPOsf was pegylated. After purification, total yield was ~30% mono-PEG-EPO with no detectable unmodified EPOsf and around 25% di-PEG-EPO. Oligomers, and unpegylated EPOsf accounted for the remaining protein. The mono-PEG-EPO pool obtained from the 150 ml fraction contained approximately 90% mono-PEG-EPO and approximately 10% di-PEG-EPO.

EXAMPLE 6: Thermostability of EPO and pegylated EPO in various formulations: Analysis by DSC (differential scanning calorimetry)

It is generally accepted that the transition temperature of thermal denaturation measured by differential scanning calorimetry is a valid indicator for the thermostability of proteins. Erythropoietin or pegylated erythropoietin (prepared according to Example 3) solutions with concentrations between 0.6 and 1.2 mg/ml were analyzed in various buffers with or without stabilizers by means of a Nano-DSC (Calorimetric Sciences Corporation, Utah, USA) at a heating rate of 2 K/min. An increase in transition temperature indicates an increase in thermal stability of the protein. The measured

temperature values should not be understood as absolute values but rather represent differences in the stability of the individual formulations relative to one another.

In order to define the optimal pH of the formulation, the pH-dependence of the thermal denaturation of pegylated erythropoietin in the range between 4 and 9 was studied. The protein samples were analyzed in 30 mM Na₂HPO₄, 30 mM sodium citrate, 30 mM borate. Figure 1 shows a plateau of maximal transition temperature between about pH 6 to about pH 9 and a sharp decrease below pH 5.5. This indicates that the optimal pH for maximal thermal stability lies above pH 5.5. (Fig. 1).

In order to investigate the effect of ionic strength, the phosphate concentration dependence of thermal denaturation was determined. Figure 2 shows that the thermal stability increases with an increase in ionic strength of the formulation.

The influence of the buffer substance was also investigated by DSC. From Figure 3 one can see that the most suitable buffers or additives for a high thermal stability are sulfate, citrate or phosphate. Glycine, which is used as a buffer in currently available formulations (see above) is not very suitable.

Figure 4 shows that sulfate is also a suitable buffer/additive at low pH (e.g. pH 6.2), whereas phosphate is less suitable at pH 6.2 compared to pH 7.5. This shows that sulfate keeps the thermal stability high, even at low pH. This finding allows a formulation at a pH between 6.0 and 6.5, without severe losses in thermal stability of erythropoietin.

EXAMPLE 7: Aggregation of EPO and peg-EPO under thermal stress: Analysis by SDS-PAGE (sodium dodecyl sulphate polyacrylamide gel electrophoresis)

In order to investigate the effect of heat stress on the erythropoietin protein, samples in different formulations were exposed to heat stress (20 min 80 °C) and analyzed by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) under reducing (with DTT in sample buffer) and non-reducing (w/o DTT in sample

buffer) conditions. This method allows the detection of covalent aggregate formation. As outlined above, aggregate formation is one of the major degradation pathways of proteins and therefore should be prevented in pharmaceutical formulations of proteins. Aggregates that are detectable in the absence of reducing agent (e.g. DTT) and not detectable in the presence of reducing agent are highly likely to be formed by incorrect disulfide bridging, an oxidation reaction, under heat stress. Figure 5 shows the pH dependency of aggregation under heat stress. This experiment clearly shows that the formation of aggregates is suppressed at a pH below 6.5. The higher the pH, the higher the amount of aggregation. Most of the aggregates that are formed can be reduced by treatment of the samples with a reducing agent during SDS-PAGE, suggesting that a great portion of the aggregates that are formed under heat stress are disulfide-bridged dimers, oligomers and higher order aggregates. Taken together, this indicates that the formation of aggregates can be prevented to a great extent by keeping the pH of the formulation at or below pH 6.5.

Figure 5: Dependency of peg-EPO aggregation on pH. Peg-EPO samples prepared according to Example 3 were subjected to heat stress (as described above) and then analyzed by SDS-PAGE. Proteins were stained with silver. Lane 1: molecular weight standard. Lane 2: pH 5. Lane 3: pH 5, reduced. Lane 4: pH 6. Lane 5: pH 6, reduced. Lane 6: pH 6.5. Lane 7: pH 6.5, reduced. Lane 8: pH 7. Lane 9: pH 7, reduced. Lane 10: peg-EPO, unstressed.

The formation of aggregates can also be prevented by the use of antioxidants. Figure 6 shows that the use of 1 mg/ml acetylcysteine as an antioxidant prevents the formation of aggregates under heat stress. Therefore, it is useful to use an antioxidant, like e.g. acetylcysteine at a low pH, e.g. pH 6.2, to prevent aggregate formation under heat stress.

Figure 6: Peg-EPO aggregation can be prevented by pH 6.2 and/or acetylcysteine. Peg-EPO samples prepared according to Example 3 were subjected to heat stress (as described above) and then analyzed by SDS-PAGE. Proteins were stained with silver.

Lane 1: peg-EPO, unstressed. Lane 2: pH 7.5, stressed. Lane 3: pH 6.2, stressed. Lane 4: pH 6.2, stressed, reduced. Lane 5: pH 7.5, 1 mg/ml acetylcysteine, stressed. Lane 6: pH 7.5, 1 mg/ml acetylcysteine, stressed, reduced.

5 **Example 8: Stability of peg-EPO in various formulations at 4, 25, 30 and 40 °C**

Pegylated EPO prepared according to Example 3 in various formulations is incubated at several temperatures. At indicated time points, samples are taken and the stability is assessed by reversed phase high performance chromatography (rpHPLC), high performance size exclusion chromatography (SEC) and sodium dodecyl sulphate
10 polyacrylamide gel electrophoresis (SDS-PAGE). Table 3 compares the stability of peg-EPO in various formulations at several temperatures. These data clearly show the superiority of the herein enclosed formulations regarding protein recovery and aggregation.

Table 3: Stability of peg-EPO in various formulations at several temperatures:

Formulation*	PegEPO (µg/ml)	% recovery after one month at				Aggregation at 40 °C detectable (+/-)
		4°C	25°C	30°C	40°C	
A	10	95	92	n.d.	66	-
B	10	93	90	n.d.	64	-
C	10	115	115	111	105	-
D	10	100	99	102	93	-
E	50	n.d.	106	99	84	+
F	50	98	100	98	89	-
G	50	101	101	101	100	-
H	50	105	103	101	102	-
I	50	103	101	104	104	-
A	100	100	99	n.d.	79	+
B	100	103	100	n.d.	77	+
C	100	103	102	103	88	-
D	100	105	106	106	98	-
E	400	98	96	89	88	+
F	400	99	97	96	93	-
G	400	98	96	100	106	-
H	400	107	108	102	97	-
I	400	104	105	98	103	-

*the formulations are:

formulation A: 10 mM sodium phosphate, 100 mM sodium chloride, pH 7.5.

formulation B: 200 mM glycine, pH 7.1 .

5 **formulation C:** 10 mM sodium phosphate, 140 mM sodium sulfate, pH 6.2.

formulation D: 10 mM sodium phosphate, 40 mM sodium sulfate, 4% (w/v) mannitol,
pH 6.2.

formulation E: 10 mM sodium phosphate, 100 mM NaCl, pH 7.0.

formulation F: 10 mM sodium phosphate, 120 mM sodium sulfate, pH 6.2

formulation G: 10 mM sodium phosphate, 40 mM sodium sulfate, 3% (w/v) mannitol,
pH 6.2.


formulation H: 10 mM sodium phosphate, 40 mM sodium sulfate, 3% (w/v) mannitol,
7.5 μ M CaCl_2 , pH 6.2.

5 **formulation I:** 50 mM arginine, 100 mM sodium sulfate, 1 mM CaCl_2 , pH 6.2.

What is claimed is:

1. A conjugate, said conjugate comprising an erythropoietin glycoprotein having at least one free amino group and having the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells and selected from the group consisting of human erythropoietin and analogs thereof which have sequence of human erythropoietin modified by the addition of from 1 to 6 glycosylation sites or a rearrangement of at least one glycosylation site; said glycoprotein being covalently linked to "n" poly(ethylene glycol) groups of the formula $-\text{CO}-(\text{CH}_2)_x-(\text{OCH}_2\text{CH}_2)_m-\text{OR}$ with the $-\text{CO}$ of each poly(ethylene glycol) group forming an amide bond with one of said amino groups; wherein R is lower alkyl; x is 2 or 3; m is from about 450 to about 900; n is from 1 to 3; and n and m are chosen so that the molecular weight of the conjugate minus the erythropoietin glycoprotein is from 20 kilodaltons to 100 kilodaltons.

2. The conjugate of claim 1, of the formula:



wherein m, n and R are as above, and P is the residue of the glycoprotein without the n amino group(s) which form amide linkage(s) with the poly(ethylene glycol) group(s).

3. The conjugate of claim 2, wherein the glycoprotein is a human erythropoietin.

4. The conjugate of claim 3, wherein the human erythropoietin glycoprotein is expressed by endogenous gene activation.

5. The conjugate of claim 3, wherein the glycoprotein has the sequence SEQ ID NO:1.

6. The conjugate of claim 5, wherein R is methyl.

7. The conjugate of claim 5, wherein m is from about 650 to about 750.
8. The conjugate of claim 5, wherein n is 1.
- 5 9. The conjugate of claim 5, wherein R is methyl; m is from about 650 to about 750; and n is 1.
10. The conjugate of claim 2, wherein the glycoprotein has the sequence of human erythropoietin modified by the addition of from 1 to 6 glycosylation sites.
- 10 11. The conjugate of claim 10, wherein the glycoprotein has the sequence of human erythropoietin modified by a modification selected from the group consisting of:
 - Asn³⁰Thr³²;
 - Asn⁵¹Thr⁵³,
 - 15 Asn⁵⁷Thr⁵⁹;
 - Asn⁶⁹;
 - Asn⁶⁹Thr⁷¹;
 - Ser⁶⁸Asn⁶⁹Thr⁷¹;
 - Val⁸⁷Asn⁸⁸Thr⁹⁰;
 - 20 Ser⁸⁷Asn⁸⁸Thr⁹⁰;
 - Ser⁸⁷Asn⁸⁸Gly⁸⁹Thr⁹⁰;
 - Ser⁸⁷Asn⁸⁸Thr⁹⁰Thr⁹²;
 - Ser⁸⁷Asn⁸⁸Thr⁹⁰Ala¹⁶²;
 - Asn⁶⁹Thr⁷¹Ser⁸⁷Asn⁸⁸Thr⁹⁰;
 - 25 Asn³⁰Thr³²Val⁸⁷Asn⁸⁸Thr⁹⁰;
 - Asn⁸⁹Ile⁹⁰Thr⁹¹;
 - Ser⁸⁷Asn⁸⁹Ile⁹⁰Thr⁹¹;
 - Asn¹³⁶Thr¹³⁸;
 - Asn¹³⁸Thr¹⁴⁰;
 - 30 Thr¹²⁵; and
 - Pro¹²⁴Thr¹²⁵.

12. The conjugate of claim 11, wherein R is methyl.

13. The conjugate of claim 11, wherein m is from about 650 to about 750.

5 14. The conjugate of claim 11, wherein n is 1.

15. The conjugate of claim 11, wherein R is methyl; m is from about 650 to about 750; and n is 1.

10 16. The conjugate of claim 10, wherein the glycoprotein has a sequence comprising the sequence of human erythropoietin and a second sequence at the carboxy terminus of the human erythropoietin sequence, wherein the second sequence contains at least one glycosylation site.

15 17. The conjugate of claim 16, wherein the second sequences comprises a sequence derived from the carboxy terminal sequence of human chorionic gonadotropin.

18. The conjugate of claim 17, wherein the glycoprotein has a sequence selected from the group consisting of:

- 20 (a) the sequence of human erythropoietin and the sequence SEQ ID NO:3 at the carboxy terminus of the human erythropoietin sequence;
- (b) the sequence in (a) modified by Ser⁸⁷ Asn⁸⁸ Thr⁹⁰; and
- (c) the sequence in (a) modified by Asn³⁰ Thr³² Val⁸⁷ Asn⁸⁸ Thr⁹⁰.

25 19. The conjugate of claim 18, wherein R is methyl.

20. The conjugate of claim 18, wherein m is from about 650 to about 750.

21. The conjugate of claim 18, wherein n is 1.

30

22. The conjugate of claim 18, wherein R is methyl; m is from about 650 to about 750; and n is 1.

23. The conjugate of claim 2, wherein the glycoprotein has the sequence of human erythropoietin modified by a rearrangement of at least one glycosylation site.

24. The conjugate of claim 23, wherein the rearrangement comprises deletion of any of the N-linked glycosylation sites in human erythropoietin and addition of an N-linked glycosylation site at position 88 of the sequence of human erythropoietin.

25. The conjugate of claim 24, wherein the glycoprotein has the sequence of human erythropoietin modified by a modification selected from the group consisting of:

Gln²⁴ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰;

Gln³⁸ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰; and

Gln⁸³ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰.

26. The conjugate of claim 25, wherein R is methyl.

27. The conjugate of claim 25, wherein m is from about 650 to about 750.

28. The conjugate of claim 25, wherein n is 1.

29. The conjugate of claim 25, wherein R is methyl; m is from about 650 to about 750; and n is 1.

30. A composition comprising conjugates, each of said conjugates comprising an erythropoietin glycoprotein having at least one free amino group and having the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells and selected from the group consisting of human erythropoietin and analogs thereof which have sequence of human erythropoietin modified by the addition of from 1 to 6 glycosylation sites or a rearrangement of at least one glycosylation site; the


glycoprotein in each said conjugate being covalently linked to "n" poly(ethylene glycol) groups of the formula $-\text{CO}-(\text{CH}_2)_x-(\text{OCH}_2\text{CH}_2)_m-\text{OR}$ with the $-\text{CO}$ of each poly(ethylene glycol) group forming an amide bond with one of said amino groups; where in each of said conjugates R is lower alkyl; x is 2 or 3; m is from about 450 to about 900; n is from 1 to 3; n and m are chosen so that the molecular weight of each conjugate minus the erythropoietin glycoprotein is from 20 kilodaltons to 100 kilodaltons; the percentage of conjugates where n is 1 is at least ninety percent.

31. The composition of claim 30 wherein the percentage of conjugates where n is 1 is at least ninety-two percent.

32. The composition of claim 31 wherein the percentage of conjugates where n is 1 is at least ninety-six percent.

33. The composition of claim 30 wherein the percentage of conjugates where n is 1 is from ninety percent to ninety-six percent.

34. The composition of claim 30, wherein each of said conjugates has the formula:



wherein m, n and R are as above, and P in each conjugate is the residue of the glycoprotein without the n amino group(s) which form amide linkage(s) with the poly(ethylene glycol) group(s).

35. The composition of claim 34 wherein the percentage of conjugates where n is 1 is at least ninety-two percent.

36. The composition of claim 35 wherein the percentage of conjugates where n is 1 is at least ninety-six percent.

37. The composition of claim 34 wherein the percentage of conjugates where n is 1 is from ninety percent to ninety-six percent.

38. The composition of claim 34, where in each conjugate the glycoprotein is a human erythropoietin.

39. The composition of claim 38, where in each conjugate the human erythropoietin glycoprotein is expressed by endogenous gene activation.

40. The composition of claim 38, where in each conjugate the glycoprotein has the sequence SEQ ID NO:1.

41. The composition of claim 40, where in each conjugate R is methyl.

42. The composition of claim 40, where in each conjugate m is from about 650 to about 750.

43. The composition of claim 40, where in each conjugate R is methyl and m is from about 650 to about 750.

44. The composition of claim 34, where in each conjugate the glycoprotein has the sequence of human erythropoietin modified by the addition of from 1 to 6 glycosylation sites.

45. The composition of claim 44, where in each conjugate the glycoprotein has the sequence of human erythropoietin modified by a modification selected from the group consisting of:

Asn³⁰Thr³²;

Asn⁵¹Thr⁵³,

Asn⁵⁷Thr⁵⁹;

Asn⁶⁹;

Asn⁶⁹Thr⁷¹;
 Ser⁶⁸Asn⁶⁹Thr⁷¹;
 Val⁸⁷Asn⁸⁸Thr⁹⁰;
 Ser⁸⁷Asn⁸⁸Thr⁹⁰;
 5 Ser⁸⁷Asn⁸⁸Gly⁸⁹Thr⁹⁰;
 Ser⁸⁷Asn⁸⁸Thr⁹⁰Thr⁹²;
 Ser⁸⁷Asn⁸⁸Thr⁹⁰Ala¹⁶²;
 Asn⁶⁹Thr⁷¹Ser⁸⁷Asn⁸⁸Thr⁹⁰;
 Asn³⁰Thr³²Val⁸⁷Asn⁸⁸Thr⁹⁰;
 10 Asn⁸⁹Ile⁹⁰Thr⁹¹;
 Ser⁸⁷Asn⁸⁹Ile⁹⁰Thr⁹¹;
 Asn¹³⁶Thr¹³⁸;
 Asn¹³⁸Thr¹⁴⁰;
 Thr¹²⁵; and
 15 Pro¹²⁴Thr¹²⁵.

46. The composition of claim 45, where in each conjugate R is methyl.

47. The composition of claim 45, where in each conjugate m is from about 650 to
 20 about 750.

48. The composition of claim 45, where in each conjugate R is methyl and m is from about 650 to about 750.

25 49. The composition of claim 44, where in each conjugate the glycoprotein has a sequence comprising the sequence of human erythropoietin and a second sequence at the carboxy terminus of the human erythropoietin sequence, wherein the second sequence contains at least one glycosylation site.

50. The composition of claim 49, where in each conjugate the second sequences comprises a sequence derived from the carboxy terminal sequence of human chorionic gonadotropin.

5 51. The composition of claim 50, where in each conjugate the glycoprotein has a sequence selected from the group consisting of:

(a) the sequence of human erythropoietin and the sequence SEQ ID NO:3 at the carboxy terminus of the human erythropoietin sequence;

(b) the sequence in (a) modified by Ser⁸⁷ Asn⁸⁸ Thr⁹⁰; and

10 (c) the sequence in (a) modified by Asn³⁰ Thr³² Val⁸⁷ Asn⁸⁸ Thr⁹⁰.

52. The composition of claim 51, where in each conjugate R is methyl.

15 53. The composition of claim 51, where in each conjugate m is from about 650 to about 750.

54. The composition of claim 51, where in each conjugate R is methyl and m is from about 650 to about 750.

20 55. The composition of claim 34, where in each conjugate the glycoprotein has the sequence of human erythropoietin modified by a rearrangement of at least one glycosylation site.

25 56. The composition of claim 55, where in each conjugate the rearrangement comprises deletion of any of the N-linked glycosylation sites in human erythropoietin and addition of an N-linked glycosylation site at position 88 of the sequence of human erythropoietin.

30 57. The composition of claim 56, where in each conjugate the glycoprotein has the sequence of human erythropoietin modified by a modification selected from the group consisting of:

Gln²⁴ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰;
Gln³⁸ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰; and
Gln⁸³ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰.

5 58. The composition of claim 57, where in each conjugate R is methyl.

59. The composition of claim 57, where in each conjugate m is from about 650 to about 750.

10 60. The composition of claim 57, where in each conjugate R is methyl and m is from about 650 to about 750.

99759

ABSTRACT OF THE INVENTION

Conjugates of erythropoietin with poly(ethylene glycol) comprise an erythropoietin glycoprotein having at least one free amino group and having the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells and selected from the group consisting of human erythropoietin and analogs thereof which have sequence of human erythropoietin modified by the addition of from 1 to 6 glycosylation sites or a rearrangement of at least one glycosylation site; said glycoprotein being covalently linked to "n" poly(ethylene glycol) groups of the formula $-\text{CO}-(\text{CH}_2)_x(\text{OCH}_2\text{CH}_2)_m-\text{OR}$ with the carbonyl of each poly(ethylene glycol) group forming an amide bond with one of said amino groups; wherein R is lower alkyl; x is 2 or 3; m is about 450 to about 900; n is from 1 to 3; and n and m are chosen so that the molecular weight of the conjugate minus the erythropoietin glycoprotein is from 20 kilodaltons to 100 kilodaltons.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6



Declaration and Power of Attorney for Patent Application

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

ERYTHROPOIETIN CONJUGATES

the specification of which

(check one)

☒ is attached hereto.

☐ was filed on _____ as

Application Serial No. _____

and was amended on _____
(if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)**Priority Claimed**

_____ (Number)	_____ (Country)	_____ (Day/Month/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No
_____ (Number)	_____ (Country)	_____ (Day/Month/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No
_____ (Number)	_____ (Country)	_____ (Day/Month/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below.

<u>60/142,254</u> (Application No.)	<u>July 2, 1999</u> (Filing Date)	<u>60/150,225</u> (Application No.)	<u>August 23, 1999</u> (Filing Date)
<u>60/151,548</u> (Application No.)	<u>August 31, 1999</u> (Filing Date)	<u>60/166,151</u> (Application No.)	<u>November 17, 1999</u> (Filing Date)

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

_____ (Application Serial No.)	_____ (Filing Date)	_____ (Status) (patented, pending, abandoned)
_____ (Application Serial No.)	_____ (Filing Date)	_____ (Status) (patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. *(list name and registration number)*

George W. Johnston	(Reg. No. 28090)	Dennis P. Tramaloni	(Reg. No. 28542)
William H. Epstein	(Reg. No. 20008)	Patricia S. Rocha-Tramaloni	(Reg. No. 31054)
Briana C. Buchholz	(Reg. No. 39123)	John P. Parise	(Reg. No. 34403)

Send Correspondence to:
George W. Johnston, Esq., Hoffmann-La Roche Inc., 340 Kingsland Street,
Nutley, New Jersey 07110-1199

Direct Telephone Calls to: (name and telephone number)

William H. Epstein (973) 235-3723


Pascal Sebastian Bailon
Full name of sole or first inventor

Inventor's signature Date

Florham Park, Morris County, New Jersey
Residence

U.S.A.
Citizenship

21 Woodbine Road, Florham Park, New Jersey 07932
Post Office Address


Full name of second joint inventor, if any

Second Inventor's signature

Date

Residence

Citizenship

Post Office Address

Full name of third joint inventor, if any

Third Inventor's signature

Date

Residence

Citizenship

Post Office Address

Full name of fourth joint inventor, if any

Fourth Inventor's signature

Date

Residence

Citizenship

Post Office Address

Full name of fifth joint inventor, if any

Fifth Inventor's signature

Date

Residence

Citizenship

Post Office Address

Title 37, Code of Federal Regulations, §1.56, duty to disclose information material to patentability provides, in part, that each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is cancelled or withdrawn from consideration, or the application becomes abandoned.

Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and

- (1) It establishes, by itself or in combination with other information, a prima facie case of unpatentability of a claim; or
- (2) It refutes, or is inconsistent with, a position the applicant takes in:
 - (i) Opposing an argument of unpatentability relied on by the Office, or
 - (ii) Asserting an argument of patentability.

1097.app
SEQUENCE LISTING

<110> Bailon, Pascal

<120> ERYTHROPOIETIN CONJUGATES

<130> 1097 nonprovisional

<140>

<141>

<150> 60/166,151

<151> 1999-11-17

<150> 60/151,548

<151> 1999-08-13

<150> 60/150,225

<151> 1999-08-23

<150> 60/142,254

<151> 1999-07-02

<160> 3

<170> PatentIn Ver. 2.1

<210> 1

<211> 165

<212> PRT

<213> Homo sapiens

<400> 1

Ala	Pro	Pro	Arg	Leu	Ile	Cys	Asp	Ser	Arg	Val	Leu	Glu	Arg	Tyr	Leu
1				5					10					15	

Leu	Glu	Ala	Lys	Glu	Ala	Glu	Asn	Ile	Thr	Thr	Gly	Cys	Ala	Glu	His
			20					25					30		

Cys	Ser	Leu	Asn	Glu	Asn	Ile	Thr	Val	Pro	Asp	Thr	Lys	Val	Asn	Phe
		35					40					45			

Tyr	Ala	Trp	Lys	Arg	Met	Glu	Val	Gly	Gln	Gln	Ala	Val	Glu	Val	Trp
	50					55					60				

Gln	Gly	Leu	Ala	Leu	Leu	Ser	Glu	Ala	Val	Leu	Arg	Gly	Gln	Ala	Leu
65					70					75					80

[illegible]

```

<210> 2
<211> 166
<212> PRT
<213> Homo sapiens

<400> 2
Ala  Pro  Pro  Arg  Leu  Ile  Cys  Asp  Ser  Arg  Val  Leu  Glu  Arg  Tyr  Leu
   1              5              10              15
Leu  Glu  Ala  Lys  Glu  Ala  Glu  Asn  Ile  Thr  Thr  Gly  Cys  Ala  Glu  His
              20              25              30
Cys  Ser  Leu  Asn  Glu  Asn  Ile  Thr  Val  Pro  Asp  Thr  Lys  Val  Asn  Phe
              35              40              45
Tyr  Ala  Trp  Lys  Arg  Met  Glu  Val  Gly  Gln  Gln  Ala  Val  Glu  Val  Trp
      50              55              60
Gln  Gly  Leu  Ala  Leu  Leu  Ser  Glu  Ala  Val  Leu  Arg  Gly  Gln  Ala  Leu
   65              70              75              80
Leu  Val  Asn  Ser  Ser  Gln  Pro  Trp  Glu  Pro  Leu  Gln  Leu  His  Val  Asp
              85              90              95
Lys  Ala  Val  Ser  Gly  Leu  Arg  Ser  Leu  Thr  Thr  Leu  Leu  Arg  Ala  Leu
              100              105              110
Gly  Ala  Gln  Lys  Glu  Ala  Ile  Ser  Pro  Pro  Asp  Ala  Ala  Ser  Ala  Ala

```

115

120

125

Pro Leu Arg Thr Ile Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val
 130 135 140

Tyr Ser Asn Phe Leu Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala
 145 150 155 160

Cys Arg Thr Gly Asp Arg
 165

<210> 3

<211> 28

<212> PRT

<213> Homo sapiens

<400> 3

Ser Ser Ser Ser Lys Ala Pro Pro Pro Ser Leu Pro Ser Pro Ser Arg
 1 5 10 15

Leu Pro Gly Pro Ser Asp Thr Pro Ile Leu Pro Gln
 20 25